SpinQuest at FNAL

Dustin Keller
University of Virginia

September Collaboration Meeting



- SpinQuest at Fermilab Goals
- Target and Intensity Frontier
- Status of Collaboration

Fermilab

Drell-Yan





- Cross section is a convolution of beam and target parton distributions

$$\frac{d^2\sigma}{dx_{\mathrm{b}}dx_{\mathrm{t}}} = \frac{4\pi\alpha^2}{x_{\mathrm{b}}x_{\mathrm{t}}s}\sum_{q\in\{u,d,s,\ldots\}} e_q^2\left[\bar{q}_{\mathrm{t}}\left(x_{\mathrm{t}}\right)q_{\mathrm{b}}\left(x_{\mathrm{b}}\right)+\bar{q}_{\mathrm{b}}\left(x_{\mathrm{b}}\right)q_{\mathrm{t}}\left(x_{\mathrm{t}}\right)\right]$$

- u-quark dominance
  $(2/3)^2$ vs. $(1/3)^2$

Beam	Sensitivity	Experiment
Hadron	Beam quarks target antiquarks	Fermilab, J-PARC RHIC (forward acpt.)
Anti-Hadron	Beam antiquarks Target quarks	J-PARC, GSI-FAIR Fermilab Collider
Meson	Beam antiquarks Target quarks	COMPASS, J-PARC

Accessing Quark Sivers TMDs

Polarized Semi-Inclusive DIS



$$A_{UT}^{SIDIS} \propto \frac{\sum_q e_q^2 f_{1T}^{\perp,q}(x) \otimes D_1^q(z)}{\sum_q e_q^2 f_1^q(x) \otimes D_1^q(z)}$$

- L-R asymmetry in hadron production
- Quark to Hadron Fragmention function
- Valence-Sea quark: Mixed

Polarized Drell-Yan



$$A_N^{DY} \propto \frac{\sum_q e_q^2 \left[f_1^q(x_1) \cdot f_{1T}^{\perp,\bar{q}}(x_2) + 1 \leftrightarrow 2 \right]}{\sum_q e_q^2 \left[f_1^q(x_1) \cdot f_1^{\bar{q}}(x_2) + 1 \leftrightarrow 2 \right]}$$

- L-R asymmetry in Drell-yan production
- **No Quark Fragmention function**
- Valence-Sea quark **Isolated**

Polarized Drell-Yan





$$A_N\left(p_{beam}+p^{\uparrow}_{target}\rightarrow DY\right)\propto\frac{N_L^{DY}-N_R^{DY}}{N_L^{DY}+N_R^{DY}}\propto\frac{f_{1T}^{\perp,\bar{u}}\left(x_t\right)}{f_1^{\bar{u}}\left(x_t\right)}$$

$$A_N\left(p_{beam}+d^{\uparrow}_{target}\rightarrow DY\right)\propto\frac{N_L^{DY}-N_R^{DY}}{N_L^{DY}+N_R^{DY}}\propto\frac{f_{1T}^{\perp,\bar{d}}\left(x_t\right)}{f_1^{\bar{d}}\left(x_t\right)}$$

- First measurement of sea quark Sivers ($\bar{u}$, $\bar{d}$)
- Sign and value
  - Result has strong implications for O.A.M. in spin puzzle
- If nonzero, “smoking gun” for Sea quark O.A.M.
- If zero, where is proton spin coming from?

SpinQuest Goals

- Separately measure Sivers function for the sea
- Measure Sign and Magnitude
- Measurement of Sivers function for gluons (J/psi SSA)

Physics Case

- Exploring the contribution of orbital angular momentum
- Interference between spin-flip and non-flip amplitudes with phase dependence
- Soft gluons

  - Gauge link required for color gauge invariance

  - Testing interplay between time-reversal symmetry and gauge symmetry

SpinQuest Goals



- Consider a nucleonic pion cloud
  $|p\rangle = |p_0\rangle + |N\pi\rangle + |\Delta\pi\rangle + \ldots$

  Pions $J^p=0^-$ Negative Parity
  Need L=1 to get proton's $J^p=½^+$

Sea quarks should carry orbital angular momentum.

Lattice QCD:



$\Delta\Sigma_q \approx 25\%$

$2\,L_q \approx 46\%$ (0%(valence)+46%(sea))

$2\,J_g \approx 25\%$

K.-F. Liu *et al* arXiv:1203.6388

QDC Gauge Invariance

- Interference between spin-flip and non-flip amplitudes w/different phases
- Soft gluons
  - "gauge links" required for color gauge invariance
  - Re-interactions are final (or initial) state … and may be process dependent!

$$f_{1T}^{\perp}\Big|_{\mathrm{SIDIS}} = -\,f_{1T}^{\perp}\Big|_{\mathrm{DY}}$$

New World Data Fit

"Interesting" example by M. Anselmino et al (2017)

Sivers function extracted from the asymmetry



New World Data Fit

"Interesting" example by M. Anselmino et al (2017)

The sign-change (QCD prediction) is tested using the DY data of the W production from RHIC





- The chi-square pdf with/without sign-change assumption applied to the DY data from RHIC which are the only available data of Sivers asymmetry from the DY process
- No solid conclusion about the sign-change as predicted by QCD
- Our SpinQuest data will have a big impact

Proton Beam at FNAL



- 120 GeV proton beam
- $\sqrt{s}$ = 15.5 GeV
- Projected Beam for E1039
  - Beam: 5×10^{12} p/spill; spill is 5 s/min
  - Protons on target per year
    - 7×10^{17}

Advantage of the Main Injector

The (very successful) past:

Fermilab E866/NuSea

- Data in 1996-1997
- ^{1}H, ^{2}H, and nuclear targets
- 800 GeV proton beam

Fermilab E906

- Data in 2010
- ^{1}H, ^{2}H, and nuclear targets
- 120 GeV proton Beam And E1039

$$\frac{d^2\sigma}{dx_1 dx_2} = \frac{4\pi\alpha^2}{9x_1x_2s} \times \sum_i e_i^2 \left[q_{ti}(x_t)\bar{q}_{bi}(x_b) + \bar{q}_{ti}(x_t)q_{bi}(x_b)\right]$$

- Cross section scales as 1/s
  - *7× that of 800 GeV beam*
- Backgrounds, primarily from J/ψ decays scale as s
  - *7× Luminosity for same detector rate as 800 GeV beam*

50× statistics!!



Fixed Target Kinematics



$$\frac{d^2\sigma}{dx_t dx_b} = \frac{4\pi\alpha^2}{9x_1x_2s}\sum e^2[q_b(x_b)\bar{q}_t(x_t) + \bar{q}_b(x_b)q(x_t)]$$

$$\approx \frac{4\pi\alpha^2}{9x_1x_2s}\sum e^2[q_b(x_b)\bar{q}_t(x_t)]$$



Fixed target kinematics favors sea-quarks from target – **a sea quark lab!**

Experimental Setup for E1039



NM4 Detector

SpinQuest Dimuon Spectrometer



120 GeV protons from the Main Injector

- 4s beam spill very 60 sec
- 19ns RF, ~10s K protons per RF bucket
- 5×10^{12} Proton On Target (POT) per spill
- Total integrated POT for E1039 (2-year): 1.4×10^{18} POT

E906 unpolarized targets: 2012-2017

- ^{1}H, ^{2}D, ^{12}C, ^{56}Fe, ^{184}W

E1039 polarized targets: 2019 – 2021+

- Polarized protons (NH_3)
- Polarized neutrons (ND_3)

E1027 polarized beam

SeaQuest Dimuon Tracking Top View



Experimental Setup for E1039

Detector Pack



SpinQuest Experimental Hall





Beam

Target area

F-Mag

K-Mag

Muon-ID

E906 Unpolarized Physics Program

- Thin targets: ~10% interaction length
  - Liquid H/D
  - Solid C, Fe, W
- Physics
  - Sea quark flavor asymmetry, dbar/ubar
  - Quark energy loss in p+A collisions, dE/dx
  - TMD and more …
- Experimental runs – 6 years
  - 2012 – commissioning
  - 2017 – completed



Preliminary Look from SeaQuest



SeaQuest Status (E906)

Main Injector beam in a slow spill—difficult to obtain good duty factor

- 1.4×10^{18} of the 5.3×10^{18} approved "live protons"
- 1.7×10^{18} of the 7×10^{18} protons with good duty factor

3.5 x 10^{17} live protons
¼ of recorded protons



Caveats:

- Rate dependence correction has a kinematic dependence
- Leading order extraction
  - NLO code tested
- Correct method -> global fit
- Large x_{beam} dbar/ubar
- . . .

Preliminary Look from SeaQuest

Flavor Asymmetry of Sea Quarks at Intermediate x

Proton vs "Neutron" targets:

$$\left.\frac{\sigma^{pd\to\mu^+\mu^-}}{\sigma^{pp\to\mu^+\mu^-}}\right|_{x_b \gg x_t} \approx \frac{1}{2}\left[1 + \frac{\bar{d}(x_t)}{\bar{u}(x_t)}\right]$$

This could lead to a very interesting physics …

J. C. Peng et al (FNAL E866/NuSea), PRD 58, 092004 (1998)



SpinQuest Projections

Projected Drell-Yan Transverse Single Spin Asymmetry

$$A_N^{DY} \propto \frac{u(x_b) \cdot f_{1T}^{\perp,\bar{u}}(x_t)}{u(x_b) \cdot \bar{u}(x_t)}$$

$$\delta A = \frac{1}{f}\frac{1}{P}\frac{1}{\sqrt{N^+ + N^-}}$$



x_2 bin	$< x_2 >$	NH_3 $(p^{\uparrow})$		ND_3 $(d^{\uparrow})$		$n^{\uparrow}$
		N	ΔA (%)	N	ΔA (%)	ΔA(%)
0.10 - 0.16	0.139	5.0×10^4	3.2	5.8×10^4	4.3	5.4
0.16 - 0.19	0.175	4.5×10^4	3.3	5.2×10^4	4.6	5.7
0.19 - 0.24	0.213	5.7×10^4	2.9	6.6×10^4	4.1	5.0
0.24 - 0.60	0.295	5.5×10^4	3.0	6.4×10^4	4.1	5.1

$$\Delta A_N = \frac{1}{f}\frac{1}{P}\frac{1}{\sqrt{N}}$$

$$f \equiv \frac{N_{p,polarizable}}{N_p + N_n} = \frac{p \times 3}{p \times (7+3) + n \times 7} = \frac{3}{17}$$

Others: Nitrogen, Helium, Target cell, Aluminum, Thin beam window, NMR coil, …

Projections of Systematics

Error estimates (non-exhaustive)

- Statistical: 3%-5% absolute error
  - Dependent on polarization, dilution, events
  - Dependent on run time
- Systematic: Mostly relative error, some absolute.
  Numbers listed hopeful upper bounds
  - Target: ~6/7% (P/D)
    - Dilution: 3%
    - Packing Fraction: 2%
    - Density: 1%
    - Polarization: 2.5%/4.5% (P/D)
    - Polarization Homogeneity: 2%
    - Uneven Decay: 3%
    - Alignment: small absolute possible
  - Beam: 2.5%
    - Relative Luminosity: 1%
    - Drifts: 2% (Absolute possible)
    - Scraping: 1%
  - Detector: 1% (Some relative, Absolute possible)



DGLAP: M. Anselmino et al arXiv:1612.06413
TMD-1: M. G. Echevarria et al arXiv:1401.5078
TMD-2: P. Sun and F. Yuan arXiv:1308.5003

Dynamic Nuclear Polarization



Successful material for DNP characterized by three measures:

1. Maximum polarization
2. Dilution factor
3. Resistance to ionizing radiation




Material	Butanol	Ammonia, NH_3	Lithium Hydride, ^{7}LiH
Dopant	Chemical	Irradiation	Irradiation
Dil. Factor (%)	13.5	17.6	25.0
Polarization (%)	90-95	90-95	90
Material	D-Butanol	D-Ammonia, ND_3	Lithium Deuteride, ^{6}LiH
Dil. Factor (%)	23.8	30.0	50.0
Polarization (%)	40	50	55
Rad. Resistance	moderate	high	very high
Comments	*Easy to produce and handle*	*Works well at 5T/1K*	*Slow polarization, but long T_1*

- Dynamic Nuclear Polarization

  - Dope target material with paramagnetic centers:

  chemical or irradiation doping to just the right density (10^{19} spins/cm^3)

  - Polarize the centers: Just stick it in a magnetic field

  - Use microwaves to transfer this polarization to nuclei:
  mutual electron-proton spin flips re-arrange the nuclear Zeeman populations to favor one spin state over the other

- Optimize so that DNP is performed at *B/T* conditions where electron t_1 is short (ms) and nuclear t_1 is long (minutes or hours)

General System



Polarized Target

Material	Dens. (g/cm^3)	Length (cm)	Interaction Length (cm)	Dilution Factor	Packing Fraction	$< P_z >$
NH_3	0.867	7.9	91.7	0.176	0.6	80%
ND_3	1.007	7.9	82.9	0.3	0.6	32%

- 3 probes over length of target.
- NMR expected to have 2-3% error for proton 4-5% for deuteron. Deuteron signal order of magnitude smaller.
- If coils moved outside cup, possible increase in uncertainty for deuteron.
- Need time to thermalize. Need 3xt1 (relaxation rate, ~10 min for proton, 1 hour for deuteron). 2-3x more error if rushed.
- Built-in error for neutron polarization from deuteron.

$$\Delta A_N = \frac{1}{f}\frac{1}{P}\frac{1}{\sqrt{N}}$$

$$f \equiv \frac{N_{p,polarizable}}{N_p + N_n} = \frac{p \times 3}{p \times (7+3) + n \times 7} = \frac{3}{17}$$

$$f \equiv \frac{N_{p,polarizable}\ \sigma_{\pi p}^{DY}}{N_p \sigma_{\pi p}^{DY} + N_n \sigma_{\pi n}^{DY}}$$







Firsts for Polarized Targets

UVA-LANL: Three completely new NMRs



NMR

UVA: Design

Insert

Microwave

CPI

UVA: Tune System and Automation



Pumps

œrlikon

14,000 providing the highest cooling power for 1K system



NIST National Institute of Standards and Technology U.S. Department of Commerce

Target material

UVA: Target Insert with longest cell at 8 cm for 5T

Fridge

UVA: Configure Fridge and Insert, Commission for Optimal running, setup with Actuator

Magnet

OXFORD INSTRUMENTS

UVA: Commisioning, Slow Controls, Quench Study, Beamline interface,...

Polarized target on the Intensity Frontier

Highest Intensity proton beam on polarized target with $4x10^{12}$ per 4s spill

- 8 cm long target cell of solid:

  NH_3 and ND_3

- Several watts of cooling power:

  14,000 m^3 /hour pumping

- 5T vertically pointing SC magnet:

  Pushing critical temp each spill

- Luminosity of around $2X10^{35}$ cm^{-2} s^{-1}

Target Insert



3 NMR coils per cell

- 8 cm long target cell of solid: NH_3 and ND_3
- Standard Insert has 3 cells
- One centering cell
- Elliptically shaped to match profile

Target Performance



Insert in LN2

95%



~2.5

~1 hr to to fill magnet can



~1hr to fill the nose after a night on standby

very stable, very little attention required



Polarized target on the Intensity Frontier



DNP Refrigerator







Cave Setup in Fermilab NM4



SpinQuest He-Liquefier

Modern He-Gas Recovery, Purification and Liquefier system

- Model QDHRR100 (2 units: 200 LPD)
- Turn-key/low maintenance system
- 135 LPD required at target for sustainable running
- **200 LPD need before transfer (fill over 60 min.)**
- LANL/UVA purchase for SpinQuest

Cryo-Platform





DNP Fridge Access



Target Cave from Upstream



Cryo-platform



Polarized Target Frontier

SpinQuest

- Cycle Time: Every 55.6 seconds
- Spill Length: 4.4 seconds
- Beam Intensity $4.0X10^{12}$ protons/sec

VS

BNL:

Energy	24 GeV
Cycle Time	3 seconds
Spill Length	1 second
Beam Intensity	2×10^{11} protons/pulse

Limiting Factors: - Fridge Cooling Power
- Heat load to SC Magnet
- Cycle Time

BNL : $4.0X10^{12}$ protons/min - 4 cm

FNAL : $4.4X10^{12}$ protons/min - 8 cm

Highest Cooling Power DNP Evaporation System:

- Running at 20 SLPM have 1.4 W of cooling power
  - For 4.4 sec receive 0.4 W of heat load from protons
  - Continuous DNP microwave heat load 0.65 W
- Super conducting magnet critical temperature 7.5 K @ 5T
- Cycle gives time to cool

Field Measurement

Accurate Field Map ✓

We achieve a high level of homogeneity around the target area & along the beam line:



High level of homogeneity in the target area



E1039



BNL target











SpinQuest Beam Collimator







Geant → COMSOL













Quench Threshold

Introduction: Quench definition

Critical surface of a LHC NbTi wire



The critical surface is defined from the temperature (T), magnetic field (B), and the surface current (J)

Magnet quench if the T, B or J lie outside the critical surface

For B = 5 T, The maximum temperature that the magnet can hold is around 7.2 K

Physics of the Quench

Approximation Strategy



Various regimes of the heat transfer from solid to LHe

First, Steady state Film boiling regime is applied

$$h(T_s, T_{He}) = a_{FB\text{-}I}(T_s - T_{He}).$$

Second, we consider the superconducting magnet as a composite material with the effective thermal parameter



Rayleigh's model consist of parallel cylinders embedded in a continuous matrix

Rayleigh's formula

$$\frac{k_{eff}}{k_m} = 1 + \frac{3\phi}{\left(\frac{k_1 - 2k_m}{k_1 - k_m}\right) - \phi + 1.569\left(\frac{k_1 - k_m}{3k_1 - 4k_m}\right)\phi^{\frac{10}{3}} + \ldots}$$

Third, we parameterize some of the unknown properties by the effective surfaces that are in direct contact with the LHe:

- Perimeter of the He void
- Insulation
- Former





Microscopic view of the cable

SpinQuest Target Manget Analysis

Electromagnetic – Thermal Analysis

Critical surface of a LHC NbTi wire





Determine Heatload

What we have currently



Maximum hot spot
around 18000 W/m^3

Former



Simulation result

Maximum temperature of
coil around 5.7 K

Historical Test

Results on BNL experiment

The maximum temperature of the coil as a function of time



Maximum Temperature profile Tmax(t) for BNL:

- 240 GeV proton
- 2e11 proton/s
- Teflon Target

Notes:

- The BNL magnet was quenched in this setup (Teflon target & 2e11 proton/s)
- The simulation results "indicate" quench -> The heat is accumulated over time
- There is an issue about numerical convergence issue for longer run that need to be fixed -> require extremely fine Mesh and time step

SpinQuest Target Magnet

The maximum temperature of the coil as a function of time



Maximum Temperature profile Tmax(t) for E1039:

- 120 GeV proton
- 1e12 proton/s
- NH3 Target

Conclusion: It is save to run at 1e12 proton/s but I recommend this intensity to be considered as the upper limit

Prep for Quench Commissioning

Before Commissioning run

- Fix the numerical convergence issue
- Overleaf documentation (collaborative LaTex editor)
- Fine tuning geometry
- Systematic study
- Install 8 temperature sensor (Carlos)
- Create Temperature prediction for those sensors as a function of beam intensity

During Commissioning run

- Compare the simulation prediction vs experiment



Status of Collaboration

Collaboration

INSTITUTION	FULL MEMBERS	AFFILIATE MEMBERS
Abilene Christian University	Donald Isenhower (PI), Michael Daugherity, Shon Watson	Haley Stien, John Marsden, Mitchell Schneller, Nathan Rowlands, Roy Salinas, Rusty Towell, Shannon McNease, Yves Ngenzi, Thomas Fitch
Argonne National Laboratory	Paul Reimer (PI), Donald Geesaman	Kawtar Hafidi, Kevin Bailey, Thomas O'Connor, Zhihong Ye, Benjamin Zeidman
Fermi National Accelerator Laboratory	Richard Tesarek (PI), Carol Johnstone, Charles Brown	
KEK	Shin'ya Sawada (PI)	Shigeru Ishimoto
Los Alamos National Laboratory	Kun Liu (SP), Mikhail Yurov, Chun-Min Jen, Ming Liu, Xuan Li, Walter Sondheim, Zhaohuizi Ji	Jan Boissevain, Melynda Brooks, Matt Durham, David Kleinjan, Sho Uemura, Cesar Da Silva, Patrick McGaughey, Andi Klein
Mississippi State University	Lamiaa El Fassi (PI)	Dipangkar Dutta
New Mexico State University	Stephen Pate (PI), Vassili Papavassiliou, Haiwang Yu, Abinash Pun, Forhad Hossain	
RIKEN	Yuji Goto (PI)	
Tokyo Institute of Technology	Kenichi Nakano (PI), Toshi-Aki Shibata	
University of Colorado, Boulder	Edward Kinney (PI)	
University of Illinois, Urbana-Champaign	Jen-Chieh Peng (PI), Yen-Chu Chen, Ching Him Leung	Naomi Makins, Daniel Jumper, Jason Dove, Mingyan Tian, Bryan Dannowitz, Randall McClellan, Shivangi Prasad
University of Michigan	Wolfgang Lorenzon (PI), Minjung Kim, Noah Wuerfel	Daniel Morton, Richard Raymond, Marshall Scott
University of New Hampshire	Karl Slifer (PI), David Ruth	Maurik Holtrop
University of Virginia	Dustin Keller (SP), Joshua Hoskins, Zulkaida Akbar, Carlos Ramirez	Donal Day, Donald Crabb, Jixie Zhang, Oscar Rondon, Liliet Diaz, Arthur Conover, Brandon Kriesten, Simonetta Liuti, Ellen Brown, Blaine Norum, Matthew Roberts
Yamagata University	Yoshiyuki Miyachi (PI), Genki Nukazuka	Takahiro Iwata, Norihiro Doshita

Current Coordinators

Co-spokespersons
K. Liu (LANL) D. Keller (UVA)

Collaboration Coordinators

Run Coordinator x 2	Physics Coordinator	Analysis Coordinator	Systems Coordinator	Shift Coordinator
R. Tesarek (FNAL)	K. Nakano (TokyoTech)	Z. Akbar (UVA)	C. Ramirez (UVA)	Y. Miyachi (Yamagata)
J. Hoskins	Z. Akbar	K. Nakano	Y. Sumo	D. Ruth

Talks Coordinator	Information Coordinator	International Coordinator	Outreach Coordinator	Service Coordinator	Backup Run Coordinator
W. Lorenzon (Michigan)	S. Pate (NMSU)	S. Sawada (KEK)	T.-A. Shibata (TokyoTech)	P. Reimer (ANL)	J. Hoskins (UVA)
V. Papavassiliou	K. Liu	T.-A. Shibata	S. Sawada	D. Keller	M. Yurov

Critical Systems Experts

Polarized Target	Wire Chambers	Hodo.	Prop. tubes	Beam Cerenkov	Slow Control	Offline Software	NIM Trigger	FPGA Trigger	L3 Trigger	DAQ	Online Software
D. Keller (UVA)	C.-M. Jen (LANL)	F. Hossain (NMSU)	Helen Ji (LANL)	R. Tesarek (FNAL)	M. Yurov (LANL)	Postdoc (NMSU)	F. Hossain (NMSU)	M. Kim (UMich)	New PD (MSU)	Y.-C. Chen (UIUC)	K. Nakano (Tokyo)
C. Ramirez (UVA)	C. Brown (FNAL)	C. Ramirez (UVA)	X. Li (LANL)		D. Ruth (UNH)	Z. Akbar (UVA)	C. Brown (FNAL)	N. Wuerfel (UMich)	L. El Fassi (MSU)	Z. Zang (LANL)	K. Liu (LANL)
J. Hoskins (UVA)	K. Nakano (Tokyo)	D. Isenhower (ACU)			J. Hoskins (UVA)					P. Reimer (ANL)	
Z. Akbar (UVA)	L. El Fassi (MSU)									H. Leung (UIUC)	
D. Ruth (UNH)	Y. Goto (RIKEN)										
G. Nukazuka (Yama)	Y. Miyachi (Yama)										
F. Hossain (NMSU)											
A. Conover (UVA)											

Summary and Outlook

Experiments	Run Time	Collision Types	Physics
E906	2012-2017	p + targets (H, D, C, Fe, W)	- dbar/ubar asymmetry - quark dE/dx
E1039	2018 – 2021+	p + pol. targets (NH_3, ND_3)	Sea-quark Sivers, TMDs
E1067(para.) DarkQuest	2017-2021+(para.) 2021+ (dedicated)	p + any targets	dark photon, dark Higgs, ALP …
E1027	202x	Pol. p-beam +	- quark Sivers - TMD, spin

Where we are Going

TMDs probed via DY at SeaQuest

Boer-Mulders functions:

- Unpolarized Drell-Yan: $d\sigma_{DY} \propto h_1^{\perp}\bar{h}_1^{\perp}\cos(2\phi)$ — E906, E1039, E1027

Sivers functions:

- Single transverse spin asymmetry in polarized Drell-Yan:

$A_N^{DY} \propto f_{1T}^{\perp}(x_q) f_{\bar{q}}(x_{\bar{q}})$ — E1039, E1027

Transversity distributions:

- Double transverse spin asymmetry in polarized Drell-Yan:

$A_{TT}^{DY} \propto h_1(x_q) h_1(x_{\bar{q}})$ — E1027

- Drell-Yan and SIDIS involve different combinations of TMDs
- Drell-Yan does not require knowledge of the fragmentation functions
- T-odd TMDs are predicted to change sign from DIS to DY

(Boer-Mulders and Sivers functions)

Remains to be tested experimentally! →COMPASS, RHIC, EIC/SeaQuest for sea quarks



Join The Effort

- https://spinquest.fnal.gov/
- http://twist.phys.virginia.edu/E1039/

Fermilab